
Reverse phase HPLC of class I HLA eluted peptide ligands



Fraction number

FIG. 1

551.0 550 550 550 550 548.8 540.2 540 540 540 540 537.2 532.3 530.8 530 530 530 530 (30 scans, baseline subtracted, smoothed) baseline subtracted, smoothed) baseline subtracted, smoothed) (20 scans, baseline subtracted, smoothed) 526.3 m/z, amu m/z, amu m/z, amu m/z, amu 525.2 525.0 520 520 520 517.2 517.2 511.3 510 510 510 510 (30 scans, (30 scans, 507.2 200 500 500 500 B\*1510 fraction 8 B\*1503 fraction 8 B\*1508 fraction 8 B\*1501 fraction 8 365 -9e5 -**6e5** -365. 2e5 165 5.0e5 1.0e6 1.6e6 8.0e5 Intensity, cps Intensity, cps Intensity, cps Intensity, cps

Ion maps of peptides eluted from various B15 class I sHLA molecules. Mapping was accomplished with a nano-spray needle and an ESI mass spectrometer. The figure shows that the same ion peak is present in 3 of 4 B15 class I.

FIG. 2

MS/MS fragmentation-sequencing of ion 517.2 from the various B15 class I sHLA molecules. This data was accomplished by completing a second nanospray of the peptides in fraction 8 from the HPLC. This demonstrates how ions can be MS ion mapped and subsequently MS/MS sequenced. There is sufficient peptide present to do multiple MS/MS fragmentation runs. There is also sufficient peptide present to facilitate a submotif on fraction 8 or further separation in the event that two peptides had mapped at 517.2 in the ion map.





FIG.



infected and uninfected cells were compared. Ion 484.72 was unique to the HIV infected cells. Ion 484.72 was subjected to MS/MS fragmentation-sequencing. We called GPRTAALGLL as the sequence of the ligand. We synthesized this peptide and found that it sHLA B\*0702 was secreted from HIV infected and uninfected cells. The ion maps of the peptides eluted from sHLA B\*0702 in generated the same MS/MS fragmentation pattern as the ligand from HIV infected cells. This MS/MS data on a synthetic ligand matches our experimental data and validates the accuracy of our sequence.

\$ 550 1 550
#
T)
TIJ.
TU
-
Con the Brain
<b>જ</b>
0
W
<del></del>
¥
B

B\*1510


1			
456789		RRIM - F	LP LP GGX SSX SSX SSX SSX SSX SSX SSX
123	 	н <mark>О</mark> ДЫ;	N E SVSXPHAP APFTGGNGX EHVASSPAL HHAPCGVSX CHGANNDPAX XHSZPAGFAX XHSZPAGFAX XHSZPAGFAX XHSZPAGFAX XHSZPAGFAX XHSZPAGFAX XHGGBNDPAX XHGGBNDPAX XHGGBNDPAX XHGGBNDPAX XHGGBNX XHGGGBNX XHGGGBNX XHGGGNX XHGGNX XHGNX XHGGNX XHG XHG XHG XHG XHG XHG XHG XHG XHG XH
	dominant	strong	
8 9	₩ I	Es;	YMVTF APAVVGYTGF PVPNVRXNYTXSX YMVCNAEEYF ALGSZAXMPF VGYVDDYQF VGYVDDYQF VGYVDDYQF VGYVDDYQF TARVXSVEY AQHPSAXRFZAY GLGZTSAEF NAXGRESSF TARVXSVEY AAGTCGVV XLHET ILGPPGSVY XLGDVNMY VMGXTNANF AVVTXZSDF ANVTXZSDF ANTTF
234567	I I I I W E Z K H	MPPG HDI AG E	FLZAMGSTY GQYVVZPTY PMFDPPZTF XAVGHSGGTYPTYPSY EPAMVXZCF XAHTECPRGY VQGPVGVZY VQGPVGSAPYGGZY TGAPVSEGGY VQXYYGSVV GQPGAPXGGZY GQPHNGXRAY AAHWHVEAY TPPTRRESY FPTDRRSZF YTGVSYXHF AQASAPDAY GQRKGAGSVF VQYYYPF ATGTAZNXNZY VVACVY PLA-N-HTY VVACVY FOARXTEY XAGFFXXEY TGATX TPAZAF
1	dominant - (	strong L L V	WDRHTXF ALGARGY XSVEY XGSCDYZARGY ADPHAPPZY XAZVZMTAY NQZHGSAEY FGXACXATSY RGACXATSY APWARGZY TGAY GQZZAVDF TGAY GQLASVSY FCSNHAY NPPAZZPNSY NPPAZZPN -O-DPPPDMZY XQAGGZY SQFGGGSQY SQFGGGSQY SQFGGGSQY SQFGGGSQY CO-DPPPDMZYAAHVPTYAAHVPTYAAHVPTYAAHVPTYAAHVPTYAAHVPTY CPLSCFT FLZAMZSTY TVXDSZTHY AQAAPFAGYFGHY ALWPZF VPHZNAY ALWPZF
6 8 2	<b>₩</b>	(Sq.	ILGPPGSVY XLGDVNMY APXVSY YMSGXYGTFAVVAZSY MPAGYNNVY XPVVPAAZTY YMIDPSGVSY
1234567	, м м м м м м м м м м м м м м м м м м м	   日   日   日   日   日   日   日	AQYAAGESFY TPHTZHDEY YMFMY WVGXXPAT DPEYVSGEZE VVACVY VVAPITTGY XAXYRRMY PXAMQXXTYTZSNAY MAAMVGVAVY LPEQPLATY PYAMGXXYYTZSNAY MAAMVGVAVY LPEQPLATY PYAMGXXYYTZSNAY MAAMVGVAVY YVAPAF YVVK-VF AAFVGAXESY YVAPAF VGYPAF VGYPAF VGYSTY SPITTHAVAF XASYIY YGYVDDTGF ZATIAGEMIAF TARVXSVEY TARVATAGEMIAF TARVXSVEY TARVXSVEY TARVXSVEY TARVXSVEY TARVATAGEMIAF TARVXSVEY TARVXSVEY TARVXSVEY TARVXSVEY TARVXSVEY TARVXSVEY TARVATAGEMIAF TARVXSVEY TARVXSVEY TARVATAGEMIAF TARVXSVEY TARVATAGENIA TARVXSVEY TA
-	dominant	strong	WDRHTXFVT XAZYEHTY NGXAMHWTY VPCGAZSY TAZXHRGY NQZHGSAEY TGAYCZDEV NGYDGFNAGY TPXGEFYZSY TYXBZTHY VXANVT VXBZTHY VXGYGEYPGFY TYXBZTHY TYXBZTHY TYXBZTHY TYXBZTHY TYXBZTHY TYXGEYPGFY TYXGYGEYPGFY TYXGYGEYPGFYPGFY TYXGY TXXGY

## Pooled Peptide Motif

P1 P2 P3 P4 P5 P6 P7 P8 P9

<u>\_</u> \_ v

 $\kappa$   $\pi$  Q  $\Xi$   $\succ$   $\times$  D  $\Xi$ 



## Narrowing search parameters using fraction motifs:

## Ovarian Carcinoma Immunoreactive Antigen

MNGRADFREP	NAEVPRPIPH	IGPDYI <b>PTEE</b>	ERRYFAECND	<b>ESFWFRSYPL</b>
<b>AATSMLITQG</b>	LISKGILSSH	PKYGSIPKLI		KLSYVKICGE
KF KKLENSPL	GEALRSGQAR	RSSPPGHYY	KSKYDSSVSG	QSSFVTSPAA
QSSFVTSPA,A	ONEMLPHYE	PIPFSSSMNE	SAPTGITDHI	YDGPDPNLEE
SPKRKNITYE	ELRNKNRESY	EVSLT@KTDP	SVRPMHERVP	- KKEVKVNKYG
DTWDE				

Scanning with whole-pooled motif revealed 4 putative epitopes.

## Ovarian Carcinoma Immunoreactive Antigen

MINGRADFREP	NAEVPRPIPH	IGPDYIPTEE	ERRYFAECND	ESFWFRSYPL
AATSMUTQG	LISKGILSSH	PKYGSIPKU	LACMGYFAG	KLSYVKTCQE
KFKKLENSPL	GEALRSGQAR	RS SPPGHYYQ	KSKYDSSVSG	QSSFVTSPAA
GSSFVTSPAA	ONIEMLPHYE	PIPFSSSMNE	SAPTGITDHI	YGGPDPNLEE
SPKRKNITYE	ELRINKINRESY	EVSLTQKTDP	SVRPMHERVP	KKEVKVNKYG
DTWDE				

Scanning with fraction 48 peptide motif revealed 1 putative epitope.

FIG. 8

Motif Data (Edman sequencing)

		1 Sec. 164	2	%	ploj	6	%	(10)	4	%	fold	(ę)		100	%	fold	7	ી છો	8	%	plot	% ∃ ⊚
Dominant						Ī	W CO	100 100 100 100 100 100 100 100 100 100					77								a zero	
2 t 6 ld							7/Edu	7707	T	T												
3.5 1010	3,					2	L COL	1 V V V	1	T												
more case						•	ALCON	2000														
prior round									T	T												
21001 1015																-						***
										T							1					-
Strong	211	4v: [	۵	53.80	257	3	riste	(e) 5 (5)	_	8.30	2.97											
25.25.694				3		8		S. 1.5										712				
rease over		3.00				=		4V/S	T							##			7		200	
niciease over										T											715	
2 70										T								1				
1/10/1/						χ.	The Control	10.4				L L	Aliema 1 2)	919181			يديد الا	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6			
2 0-2 5 fold																					363	
2.0-2.3 1010																						-
nciedade over									Ī	T												
200												į						==				
Trace			٥	3.60	1 70	0	10 6 G	15.e. ft	×	11 00	1 63		18.500 P	Q (60)			- 基		C	11.40	1.93	
1 50-2 C fold			3						S	00.9	1.68	T.	2,926) 4	8		1404	4.		z	5.70	1.55	
increase over							_		>	5.30	1.99	Ę	1180	10,00%					2.0			
orior round		; ;				Ī										-						
	The second second						A 15				#	のなるのである				4						***************************************

FIG. 11

TU

FIG. 12

DESIGN OF HLA LIGAND/MOTIF DATABASE




FIG. 13

Entity-Relationship (ER) Diagram for HLA Ligand/Motif Database




FIG. 14

UML Diagram for HLA Ligand/Motif Database

